

Exhibit Q

Threatened and Endangered Species

Mist Resiliency Project
March 2024

Prepared for



Northwest Natural Gas

Prepared by



This page intentionally left blank

Table of Contents

1.0 Introduction 1

2.0 Identification of Species – OAR 345-021-0010(1)(q)(A)..... 1

 2.1 Desktop Review 2

 2.2 Field Surveys..... 4

 2.2.1 Botanical Survey 5

 2.2.2 General Biological and Habitat Categorization Survey..... 5

3.0 Occurrence and Potential Adverse Effects – OAR 345-021-0010(1)(q)(B) 5

 3.1 Wildlife 5

 3.1.1 Columbian White-tailed Deer (Lower Columbia River Population Only)..... 5

 3.2 Fish..... 6

 3.2.1 Coho Salmon (Oregon Coast ESU and Lower Columbia River ESU Only)..... 6

 3.2.2 Chinook Salmon (Lower Columbia River ESU Only)..... 7

 3.3 Plants..... 7

 3.3.1 Tall Bugbane 8

 3.3.2 Willamette Valley Larkspur..... 9

 3.3.3 Peacock Larkspur10

 3.3.4 Coast Range Fawn-lily10

 3.3.5 Queen-of-the-Forest.....11

 3.3.6 Howell’s Montia.....11

 3.3.7 Saddle Mountain Saxifrage12

 3.3.8 Meadow Checkermallow12

 3.3.9 Bristly-stemmed Sidalcea.....12

 3.3.10 Nelson’s Sidalcea.....13

 3.3.11 Oregon Sullivantia.....14

4.0 Avoidance and Mitigation – OAR 345-021-0010(1)(q)(C)..... 14

 4.1 Wildlife 14

 4.1.1 Fish 14

 4.2 Plants.....15

 4.2.1 Howell’s Montia.....15

5.0 Protection and Conservation Program Compliance/Impacts – OAR 345-021-0010(1)(q)(D).....15

6.0 Potential Impacts to Plants, Including Mitigation Measures – OAR 345-021-0010(1)(q)(E)16

6.1 Species Found Not to Occur16

6.2 Species Found or Previously Known to Occur17

7.0 Potential Impacts to Animals, Including Mitigation Measures – OAR 345-021-0010(1)(q)(F)17

8.0 Monitoring – OAR 345-021-0010(1)(q)(G).....17

9.0 References17

List of Tables

Table Q-1. Federal and State Listed and Candidate Species with the Potential to Occur within the Analysis Area 3

Table Q-2. Plant Blooming Period, Occurrence, and Likelihood of Adverse Effects 8

List of Figures

- Figure Q-1. Threatened and Endangered Species Analysis Area
- Figure Q-2. Surveyed Area: Threatened and Endangered Plant Species
- Figure Q-3. Endangered, Threatened, and Candidate Species Occurrences (**Confidential**)
- Figure Q-4. Howell’s Montia Observed During Surveys (**Confidential**)

Acronyms and Abbreviations

| | |
|---------|--|
| BMP | best management practice |
| ESU | Evolutionarily Significant Unit |
| HDD | Horizontal directional drill |
| NWN | Northwest Natural Gas |
| OAR | Oregon Administrative Rules |
| ODA | Oregon Department of Agriculture |
| ODFW | Oregon Department of Fish and Wildlife |
| ORBIC | Oregon Biodiversity Information Center |
| ORS | Oregon Revised Statutes |
| Project | Mist Resiliency Project |
| RFA | Request for Amendment |
| USFWS | U.S. Fish and Wildlife Service |

This page intentionally left blank

1.0 Introduction

Northwest Natural Gas (NWN), the Certificate Holder, proposes to amend the Site Certificate for its underground natural gas storage facility at the Mist Resiliency Project (Project) in Columbia County, Oregon. Exhibit Q provides information pertaining to threatened, endangered, and candidate plant and animal species that may be affected by the Project, as required to meet the submittal requirements in Oregon Administrative Rules (OAR) 345-021-0010(1)(q) paragraphs (A) through (G). This exhibit demonstrates that the Project can comply with the approval requirements found in OAR 345-022-0070:

To issue a site certificate, the Council, after consultation with appropriate state agencies, must find that:

(1) For plant species that the Oregon Department of Agriculture has listed as threatened or endangered under ORS 564.105(2), the design, construction and operation of the proposed facility, taking into account mitigation:

(a) Are consistent with the protection and conservation program, if any, that the Oregon Department of Agriculture has adopted under ORS 564.105(3); or

(b) If the Oregon Department of Agriculture has not adopted a protection and conservation program, are not likely to cause a significant reduction in the likelihood of survival or recovery of the species; and

(2) For wildlife species that the Oregon Fish and Wildlife Commission has listed as threatened or endangered under ORS 496.172(2), the design, construction and operation of the proposed facility, taking into account mitigation, are not likely to cause a significant reduction in the likelihood of survival or recovery of the species.

The Analysis Area is the area for which NWN must describe the impacts of the proposed Project changes in this Request for Amendment (RFA) 13. The Analysis Area for Exhibit Q is defined as the Site Boundary, as surface facilities related to an underground gas storage reservoir, transmission lines, and pipelines are exempted from the 5-mile study area applicable to other energy facilities pursuant to OAR 345-001-0010(35)(f) and (g). This Analysis Area applies to all species in Exhibit Q. The Site Boundary is defined in the Project Description section of this RFA 13 that reflects the information required by OAR 345-021-0010(1)(a) and (b). The Analysis Area is shown in Figure Q-1.

2.0 Identification of Species – OAR 345-021-0010(1)(q)(A)

OAR 345-021-0010(1)(q) Information about threatened and endangered plant and animal species that may be affected by the proposed facility, providing evidence to support a finding by the Council as required by OAR 345-022-0070. The applicant must include:

(A) Based on appropriate literature and field study, identification of all threatened or endangered species listed under ORS 496.172(2) and ORS 564.105(2) that may be affected by the proposed facility;

In compliance with OAR 345-021-0010(1)(q)(A), NWN identified all threatened and endangered species listed under Oregon Revised Statutes (ORS) 496.172(2) (state threatened and endangered wildlife species), ORS 564.105(2) (state threatened and endangered plant species), and 16 United States Code §1533 (federal Endangered Species Act) that may be affected by the Project. Federal and state candidate and species proposed for listing were also identified due to their potential to become listed during the site certificate amendment process.

2.1 Desktop Review

NWN used a variety of sources to identify state and federal threatened, endangered, candidate, and proposed species that may be affected by the Project. Data collection has been ongoing in the area since 2013 with updates to account for changes to species status and changes to the Project.

To identify listed, candidate, and proposed species with the potential to occur within the Analysis Area, NWN requested known occurrence locations from ORBIC within the Analysis Area and vicinity. Occurrence locations outside of the Analysis Area were requested to ensure all species with the potential to occur within the Analysis Area were included in this analysis, not just species that are known to occur within the Analysis Area. Data requested covers portions of Columbia County, Oregon.

The ODFW (ODFW 2023a, ODFW 2023b), Oregon Department of Agriculture (ODA; ODA 2023), ORBIC (ORBIC 2019, ORBIC 2023a), USFWS (USFWS 1983, USFWS 2016, USFWS 2023) and National Oceanic and Atmospheric Administration, National Marine Fisheries Service (NOAA Fisheries 2022a, NOAA Fisheries 2023) websites were queried for additional literature on listed species. The StreamNet (2024) and ODFW (2023b) databases were searched for listed fish species within 5 miles of the Analysis Area. Finally, several botanical resources were reviewed to identify proposed, candidate, threatened, and endangered plant species that could occur near the Analysis Area (Burke Museum Herbarium 2022a, CPNWH 2023, ODA 2022, Oregon Flora Project 2023, WNHP 2022). The results of the literature review were used to generate a list of state and federal listed and candidate species with the potential to occur in the Analysis Area.

Once a preliminary species list was developed through the initial desktop review described above, aerial photo imagery of the habitat in the Analysis Area was compared to the species' habitat requirements to guide species-specific and general field survey efforts and inform preliminary desktop siting. Data sources analyzed included aerial photography (Google Earth 2023), National Wetland Inventory, U.S. Geological Survey National Hydrography Dataset, Natural Resources Conservation Service soils, and the Northwest Regional Gap Analysis Project data.

Based on the review of existing data, 15 species or populations listed as federally or state threatened, endangered, or candidate were identified as having the potential to occur within the Analysis Area (Table Q-1). These included one mammal, 11 vascular plants, and three fish. No

species with a status of proposed for listing were identified as having the potential to occur within the Analysis Area.

Several species initially considered for inclusion in Exhibit Q were excluded, as they are not known or expected to occur within the Analysis Area. Species excluded because they are very unlikely to occur within the Analysis Area due to lack of suitable habitat include pink sandverbena (*Abronia umbellata* var. *brevifolia*), Fender’s blue butterfly (*Icaricia icarioides fender*), Taylor’s checkerspot butterfly (*Euphydryas editha taylor*), Oregon spotted frog (*Rana pretiosa*), Pacific marten (*Martes caurina*), northern spotted owl (*Strix occidentalis caurina*; ORBIC 2023b)¹, marbled murrelet (*Brachyramphus marmoratus*; Davis et al. 2011, ORBIC 2023b), and streaked horned lark (*Eremophila alpestris strigata*; ORBIC 2023b). Species considered because their range includes Columbia County but excluded because the Project is outside the current range of the species include: Snake River Chinook salmon (fall-run, spring/summer-run), Upper Willamette River Chinook salmon, Lower Columbia River steelhead trout, Middle Columbia River steelhead trout, Upper Willamette River steelhead trout, Columbia River chum salmon (*Oncorhynchus keta*), eulachon (*Thaleichthys pacificus*), bull trout (*Salvelinus confluentus*), and green sturgeon (*Acipenser medirostris*, southern distinct population segment; NOAA Fisheries 2022a, NOAA Fisheries 2022b, ORBIC 2023a, ORBIC 2023b, StreamNet 2024).

Table Q-1. Federal and State Listed and Candidate Species with the Potential to Occur within the Analysis Area

| Species ¹ | Federal Status | State Status | Occurrence within Analysis Area ^{2,3} |
|---|----------------|--------------|--|
| Mammals | | | |
| Columbian white-tailed deer <i>Odocoileus virginianus leucurus</i> | T | T | No |
| Fish | | | |
| Coho salmon (Oregon Coast Evolutionarily Significant Unit [ESU]) <i>Oncorhynchus kisutch</i> | T | S | Yes |
| Coho salmon (Lower Columbia River ESU) <i>Oncorhynchus kisutch</i> | T | E | Yes |
| Chinook salmon (Lower Columbia River ESU, spring run and fall run) <i>Oncorhynchus tshawytscha</i> | T | SC | Yes |
| Vascular Plants | | | |
| Tall bugbane <i>Actaea elata</i> var. <i>elata</i> (syn. <i>Cimicifuga</i>) | — | C | No |

¹ ORBIC identifies a single northern spotted owl territory approximately 4 miles from the Site Boundary, which is approximately three times the territorial home range radius of the species. The lack of habitat combined with the distance from the Site Boundary to this spotted owl territory support the species’ exclusion.

| Species ¹ | Federal Status | State Status | Occurrence within Analysis Area ^{2,3} |
|--|----------------|--------------|--|
| Willamette Valley larkspur <i>Delphinium oregonum</i> | SOC | C | No |
| Peacock larkspur <i>Delphinium pavonaceum</i> (syn. <i>Delphinium menziesii</i> ssp. <i>pallidum</i>) | SOC | E | No |
| Coast Range fawn-lily <i>Erythronium elegans</i> | SOC | T | No |
| Queen-of-the-forest <i>Filipendula occidentalis</i> | — | C | No |
| Howell's montia <i>Montia howellii</i> | — | C | Yes |
| Saddle Mt. saxifrage <i>Saxifraga hitchcockiana</i> (syn. <i>Micranthes hitchcockiana</i>) | SOC | C | No |
| Meadow checkermallow <i>Sidalcea campestris</i> | — | C | No |
| Bristly-stemmed sidalcea <i>Sidalcea hirtipes</i> | — | C | No |
| Nelson's sidalcea <i>Sidalcea nelsoniana</i> | DL | T | No |
| Oregon sullivantia <i>Sullivantia oregana</i> | SOC | C | No |
| <p>Federal Status Definitions: T = Threatened, E = Endangered, C = Candidate, SOC = Species of Concern, DL = Delisted</p> <p>Oregon Status Definitions: T = Threatened, E = Endangered, C = Candidate, SC = Sensitive Critical S = Sensitive</p> <p>1. Species shown include only those that are listed as threatened, endangered, or candidate species on Oregon and/or federal lists. Other Oregon or federal designation categories are only shown for species that fit one of the primary listing categories. Other Oregon sensitive species are addressed in Exhibit P.</p> <p>2. Plant occurrence is based on documented field surveys combined with ORBIC occurrence data. Mammal occurrence is based on ORBIC data combined with observational data during general biological/habitat categorization surveys.</p> <p>3. The Analysis Area for all species is the proposed Site Boundary.</p> | | | |

2.2 Field Surveys

NWN conducted two primary field survey efforts to evaluate potential presence of state or federally listed or candidate species: a general biological and habitat categorization survey and a botanical survey.

2.2.1 Botanical Survey

NWN conducted field surveys for listed and candidate plant species in 2022 and 2023 (Figure Q-2). Surveyors used the intuitive controlled transect method to document the presence or absence of target plant species. Surveys were conducted within the Site Boundary in suitable habitat and when an area with high potential for target plants was encountered (e.g., wetland). In 2022, field surveys were conducted during the appropriate time of year to capture target species during blooming or fruiting. In 2023, surveys of additional areas were conducted in September, outside of target species blooming periods. Areas were characterized as having potentially suitable habitat for a number of species. These additional areas will be surveyed in full prior to construction. For additional details on botanical survey methods, see Exhibit P, Attachment P-1.

2.2.2 General Biological and Habitat Categorization Survey

NWN conducted wildlife and habitat categorization surveys in 2022. Transect surveys were performed to characterize habitat throughout the Site Boundary, with simultaneous searches for special status wildlife species and special habitats. No proposed, candidate, threatened, or endangered wildlife species were observed. For additional details on the general wildlife and habitat categorization survey methods, see Exhibit P, Attachment P-1.

3.0 Occurrence and Potential Adverse Effects – OAR 345-021-0010(1)(q)(B)

(B) For each species identified under (A), a description of the nature, extent, locations and timing of its occurrence in the analysis area and how the facility might adversely affect it;

In compliance with OAR 345-021-0010(1)(q)(B), this section provides a description of the nature, extent, location, and timing of occurrence in the Analysis Area of each species identified in Table Q-1 and describes how the Project might adversely affect each species. Potential adverse effects to each species that will result from the construction of the Project are based on each species' potential for occurrence within the Analysis Area.

3.1 Wildlife

3.1.1 Columbian White-tailed Deer (Lower Columbia River Population Only)

Threatened (Federal); Threatened (State)

The Lower Columbia River population of the Columbian white-tailed deer is federally listed as Threatened and is Sensitive Critical in the state of Oregon. The Columbian white-tailed deer occurs in two distinct populations separated by 200 miles of unsuitable or discontinuous habitat: (1) the Lower Columbia River population, which occurs on both banks of the river in Clatsop and Columbia

counties, Oregon, and Wahkiakum and Cowlitz counties, Washington; and (2) the Douglas County, Oregon, population.

The Columbian white-tailed deer prefers wet prairie and lightly wooded bottomlands or "tidelands" along streams and rivers; woodlands are particularly attractive when interspersed with grasslands and pastures. Habitat information is based on the Columbian White-tailed Deer Recovery Plan (USFWS 1983) and most recent 5-year review (USFWS 2016). Along the Columbia River, Sitka spruce, dogwood, cottonwood, red alder, and willow dominate the woodland vegetation. Plants commonly found in suitable adjacent grass/forb fields include fescue, orchard grass, clover, bluegrass, velvetgrass, buttercup, and ryegrass. Reed canary grass and water foxtail are common invasive species in wet sites, and species of *Rubus*, *Juncus*, *Carex*, *Rosa*, *Sambucus*, and *Symphoricarpos* are commonly used as food and cover. Historically, Columbian white-tailed deer evolved as a prairie edge/woodland-associated species and were not confined to river valleys. Breeding occurs from mid-September through late February, with a peak in November (USFWS 2016). Fawns are born in late summer (June 1–July 31) after an approximately 200-day gestation period (USFWS 2016).

Columbian white-tailed deer were not observed during the habitat categorization field survey on June 7, 8, and 9, 2022. As the Analysis Area is outside of the known range for the Columbian white-tailed deer (Columbia County 2011, ORBIC 2023b, ODFW 2017, ODFW 2023a, UFWS 2016), they are not expected to occur within the Analysis Area. However, due to the mobility of this species there is potential for individuals to occur incidentally within suitable habitat outside of their known range. Thus, while unlikely, Columbian white-tailed deer could occur incidentally within the northernmost portion of the Analysis Area, which is located just south of U.S. Highway 30 (US-30) and west of Clatskanie.

For the purposes of this Project, suitable habitat was defined as the following habitat types within the Analysis Area and species' range: westside riparian, westside lowlands conifer-hardwood forest, irrigated pasture and hay meadows. Within the Analysis Area a total of 230.9 acres of suitable habitat occurs: westside riparian (30.6 acres), westside lowlands conifer-hardwood forest (4,637.5 acres), irrigated pasture and hay meadows (0.24 acre).

Because the Analysis Area is outside of the known range for the Columbian white-tailed deer, adverse effects to this species are not expected as a result of the Project.

3.2 Fish

Two fish species that are federally or state listed or are a candidate or proposed for federal or state listing were found to have the potential to occur within the Analysis Area based on desktop review.

3.2.1 Coho Salmon (Oregon Coast ESU and Lower Columbia River ESU Only)

The Oregon Coast ESU of coho salmon is federally listed as Threatened and is Sensitive in the state of Oregon; the Lower Columbia River ESU is federally listed as Threatened and is listed as Endangered in the state of Oregon. Mature coho salmon typically enter freshwater to spawn from

late summer to late autumn, and spawning occurs between November and January. Coho salmon spawn in small tributaries with moderate to low gradient stream reaches. After spawning, eggs typically hatch within 45 to 140 days, depending on water temperature, with longer incubation in colder water (ODFW 2007). Fry emerge between early spring to early summer, and freshwater rearing occurs until the following spring when the juveniles undergo smoltification and migrate to salt water (ODFW 2010). Juvenile coho salmon migrate to the ocean as smolts in the spring, typically from late April, May, and early June. As smolts, coho may be present in estuaries for up to a month during their migration to the ocean (ODFW 2007).

Coho salmon are mapped to occur in Lindgren Creek and the Nehalem River (ODFW 2024, StreamNet 2024). The Project will cross Lindgren Creek, which is mapped as spawning habitat for this species. Lindgren Creek will be crossed via horizontal directional drill (HDD), avoiding stream impacts. Therefore, no adverse effects to coho salmon are anticipated as a result of the Project.

3.2.2 Chinook Salmon (Lower Columbia River ESU Only)

The Lower Columbia River ESU of Chinook salmon is federally listed as Threatened and is Sensitive Critical in the state of Oregon. Chinook salmon have varied adult and juvenile life histories. The Lower Columbia River ESU of Chinook represents both spring-run and fall-run populations. Spring-run Chinook salmon spend a longer rearing period in freshwater than fall-run Chinook salmon. Spring-run Chinook salmon enter freshwater in March and April and spawn in August and September; their fry emerge in winter or early spring, then rear for a year until migrating downstream the following spring. Fall-run Chinook enter freshwater in mid-August and spawn within a few weeks; their fry emerge in the spring and migrate downstream after as little as a month of freshwater rearing (ODFW 2010).

Within the Analysis Area the Nehalem River is mapped as spawning habitat for spring-run and fall-run Chinook salmon, and Lindgren Creek is mapped as spawning habitat for the fall-run (ODFW 2024). The Project will cross Lindgren Creek via HDD, avoiding stream impacts. Therefore, no adverse effects to Chinook salmon are anticipated as a result of the Project.

3.3 Plants

Eleven plant species that are federally or state listed or are a candidate or proposed for federal or state listing were found to have the potential to occur within the Analysis Area based on desktop review. Only one of these, Howell's montia (state Candidate species), was found to occur within the Site Boundary during Project surveys (Table Q-2).

Table Q-2. Plant Blooming Period, Occurrence, and Likelihood of Adverse Effects

| Species | Blooming Period ¹ | Potential for Occurrence within Analysis Area ¹ | | | | Potential Adverse Effects ² |
|----------------------------|---|--|-------------------------------|----------------|--------------------------|--|
| | | Potential Habitat within Site Boundary | ORBIC Records (Analysis Area) | County Records | Observed during Surveys? | |
| Tall bugbane | late May–early August | Yes | No | No | No | No |
| Willamette Valley larkspur | May–July | Yes | No | No | No | No |
| Peacock larkspur | April–June | Yes | No | No | No | No |
| Coast Range fawn-lily | May–June | Yes | No | No | No | No |
| Queen-of-the-forest | late May–August | Yes | No | No | No | No |
| Howell's montia | Mar–May | Yes | Yes | Yes | Yes | Yes |
| Saddle Mt. saxifrage | late May–July | Yes | No | No | No | No |
| Meadow checkermallow | May–July | Yes | No | No | No | No |
| Bristly-stemmed sidalcea | (late April) early June–August (early November) | Yes | Yes | No | No | No |
| Nelson's sidalcea | late May–mid July (September) | Yes | Yes | No | No | No |
| Oregon sullivantia | late May–August | Yes | No | Yes | No | No |

1. Peak blooming period with full blooming period in parentheses; Sources: Burke Museum Herbarium 2022a, Burke Museum Herbarium 2022b, ORBIC 2023, Oregon Flora Project 2023

2. Potential for adverse effects not considering avoidance, minimization, and mitigation measures.

3.3.1 Tall Bugbane

No Status (Federal); Candidate (State)

Tall bugbane is a state Candidate species, but it has no special status at the federal level. This coarse perennial herb grows 3–7 feet in height and blooms between late May and early August (Oregon Flora Project 2023). Tall bugbane is known to occur west of the Cascade Mountains from southwestern British Columbia through Oregon including Columbia County (Oregon Flora Project 2023). Tall bugbane grows along the margins of mature Douglas fir (*Pseudotsuga menziesii*) and bigleaf maple (*Acer macrophyllum*) forests. The primary threat is habitat degradation or loss due to

timber management practices with close spacing of newly planted trees quickly negating the initial benefit of opening up the canopy through timber harvest (Kaye and Cramer 2003, WNHP 2014). Other threats include competition from invasive weedy species, residential development, collecting by herbalists, and trampling of plants as a result of recreational use (WNHP 2014).

Tall bugbane was not observed during Project surveys, and the Oregon Flora Project (2023) atlas and ORBIC (2023b) do not include reported observations within the Analysis Area. However, because suitable habitat is present within the Analysis Area and it has been identified in the vicinity, it was included as a species with the potential to occur. In 2023, surveys of additional areas were conducted in September, outside of the tall bugbane blooming period. Potentially suitable habitat was identified. These additional areas will be surveyed in full prior to construction and if any tall bugbane are identified they will be avoided. Therefore, the species is not expected to be adversely affected by the Project.

3.3.2 Willamette Valley Larkspur

Species of Concern (Federal); Candidate (State)

Willamette Valley larkspur is a state Candidate species and a federal Species of Concern. This slender perennial herb grows 31 inches in height and blooms between May and July (Oregon Flora Project 2023). Willamette Valley larkspur is endemic to the Willamette Valley and high elevation peaks in the northern Coast Range. At low elevation sites the Willamette Valley larkspur is most commonly found in wet prairies with shrubby or Oregon ash (*Fraxinus latifolia*) overstory, but it also occurs along roadsides and fencerows and on dry oak woodlands, open hillsides, and well-drained native prairies. At high elevation sites, it occurs on open and moderately moist slopes. The primary threat to this species is continued loss of habitat to development, herbicides, disturbance associated with road maintenance, successional encroachment, and habitat invasion by exotic species (USFWS 2010).

The Willamette Valley larkspur was not observed during Project surveys, and the Oregon Flora Project (2023) atlas and ORBIC (2023b) do not include reported observations within the Analysis Area. An occurrence of an isolated population near the west bank of the Columba River in Cowlitz County, Washington, and far from other known populations suggested the potential for this species to occur in unforeseen locations, and suitable habitat exists within the Analysis Area. Therefore, it was included as a species with the potential to occur in the Analysis Area. In 2023, surveys of additional areas were conducted in September, outside of the Willamette Valley larkspur blooming period. Potentially suitable habitat for Willamette Valley larkspur was identified. These additional areas will be surveyed in full prior to construction and if any Willamette Valley larkspur are identified they will be avoided. Therefore, the species is not expected to be adversely affected by the Project.

3.3.3 Peacock Larkspur

Species of Concern (Federal); Endangered (State)

Peacock larkspur is a state Endangered species and a federal Species of Concern. This slender perennial herb grows 35 inches in height and blooms between April through June (Oregon Flora Project 2023). Peacock larkspur is endemic to the Willamette Valley of Oregon. Peacock larkspur occurs in native wet prairies, Oregon ash (*Fraxinus latifolia*) and Oregon white oak (*Quercus garryana*) woodlands, and roadsides and fences rows where suitable moisture and soil conditions are present. Peacock larkspur is most commonly found on silty soils in the Willamette River floodplain. The primary threat to this species is continued loss of habitat to development, herbicides, disturbance associated with road maintenance, successional encroachment, and habitat invasion by exotic species (ODA 2022).

Peacock larkspur was not observed during Project surveys, and the Oregon Flora Project (2023) atlas and ORBIC (2023b) do not include reported observations within the Analysis Area. The species is primarily found within Benton, Marion, and Polk counties, its largest occurrences located in William L. Finley National Wildlife Refuge in Benton County, Washington. Peacock larkspur does occur in Multnomah and Clackamas counties far from other known populations, suggesting the potential for this species to occur in unforeseen locations, and suitable habitat exists within the Analysis Area. Therefore, it was included as a species with the potential to occur in the Analysis Area. However, as the species was not observed during Project surveys, it is not expected to occur within the Analysis Area and is not expected to be adversely affected by the Project.

3.3.4 Coast Range Fawn-lily

Species of Concern (Federal); Threatened (State)

Coast Range fawn-lily is a state Threatened species and a federal Species of Concern. This wildflower grows 5–12 inches in height and blooms between May and June (Oregon Flora Project 2023). Coast Range fawn-lily is known to occur in the northern Oregon Coast Range and the Olympic Mountains of Washington where it grows at high elevation on open sites including rocky slopes and cliffs, bog edges, meadows, and rocky balds (Oregon Flora Project 2023). The primary threats are plant and habitat destruction due to timber harvest, herbivory and grazing, fungal infection, and plant collection (Guerrant 1999 as cited in Center for Plant Conservation 2020).

The Coast Range fawn-lily was not observed during Project surveys, and the Oregon Flora Project (2023) atlas and ORBIC (2023b) do not include reported observations within the Analysis Area. The Coast Range fawn-lily's known distribution is primarily restricted to high elevation sites above the Analysis Area. However, the Oregon Flora Project (2023) atlas identifies a record as low as 1,401 feet elevation, an elevation that occurs within the Analysis Area. As there was some potential for suitable habitat within the Analysis Area and the species is known from neighboring counties, it was included as a species with the potential to occur. Nevertheless, as the species was not observed during Project surveys, it is not expected to occur within the Analysis Area and is not expected to be adversely affected by the Project.

3.3.5 Queen-of-the-Forest

No Status (Federal); Candidate (State)

Queen-of-the-forest is a state Candidate species. This erect perennial herb grows 3–7 feet in height and blooms between late May and August (Oregon Flora Project 2023). Queen-of-the-forest is endemic to northwestern Oregon and southwestern Washington where it occurs on high elevation Coast Range peaks and coastal river systems. The species grows in damp sites including riverbanks, rock crevices and seeps, damp salmonberry (*Rubus spectabilis*) shrublands, and moist rock cliffs (Oregon Flora Project 2023). The primary threats are timber management activities that lead to alterations in hydrology and increased solar radiation due to canopy removal, road and bridge construction, exposure to herbicides, and plant collection (Meinke 1982, NatureServe 2023).

Queen-of-the-forest was not observed during Project surveys, and the Oregon Flora Project (2023) atlas and ORBIC (2023b) do not include reported observations within the Analysis Area. The queen-of-the-forest's known distribution is west of Columbia County and the Analysis Area. As there was some potential for suitable habitat within the Analysis Area and the species is known from neighboring counties, it was included as a species with the potential to occur. However, as the species was not observed during Project surveys, it is not expected to occur within the Analysis Area and is not expected to be adversely affected by the Project.

3.3.6 Howell's Montia

No Status (Federal); Candidate (State)

Howell's montia is a state Candidate species, but it has no special status at the federal level. This low-growing annual herb has spreading stems 1–2 inches in length and blooms between March and May (Burke Museum Herbarium 2022a, Gisler 2004). Howell's montia occurs west of the Cascades from British Columbia to California. Howell's montia occurs in sparsely vegetated moist to seasonally wet areas such as river and pond edges, cattle and game trails, open fields, vernal pools, seeps, and wet prairies (Renner et al. 2012). Threats to the species include timber harvest, road construction and maintenance, vehicles, and competition (Gisler 2004, Renner et al. 2012).

During 2022 surveys, a single population consisting of approximately 2,700 plants growing densely within the two-track road matrix comprising approximately 800-square feet (Figure Q-4; Exhibit P, Attachment P-1). Two additional Howell's montia sub-populations consisting of one plant and eight dispersed plants (35 square foot area) were 15 and 40 feet further north along the mainline road's eastern edge (Figure Q-4; Exhibit P, Attachment P-1). Impacts to this species would be avoided by identifying and marking individuals prior to Project construction and operation.

In 2023, surveys of additional areas were conducted in September, outside of the Howell's montia blooming period. Potentially suitable habitat for Howell's montia was identified. These additional areas will be surveyed in full prior to construction and if any Howell's montia are identified they will be avoided. Therefore, the species is not expected to be adversely affected by the Project.

3.3.7 Saddle Mountain Saxifrage

Species of Concern (Federal); Candidate (State)

Saddle Mountain saxifrage is a state Candidate species and a federal Species of Concern. This perennial herb has flower stalks that grow 4–10 inches tall, and it blooms between late May and July (Oregon Flora Project 2023). Saddle Mountain saxifrage is endemic to higher peaks in the north Oregon Coast Range. The species grows on grassy balds; on thin, rocky soils; and in rock crevices (Oregon Flora Project 2023). The primary threats to the species are timber harvest and grazing (Meinke 1982).

Saddle Mountain saxifrage was not observed during Project surveys, and the Oregon Flora Project (2023) atlas and ORBIC (2023b) do not include records from Columbia County. As there was some potential for suitable habitat within the Analysis Area and the species is known from neighboring counties, it was included as a species with the potential to occur. However, as the species was not observed during Project surveys, it is not expected to occur within the Analysis Area and is not expected to be adversely affected by the Project.

3.3.8 Meadow Checkermallow

No Status (Federal); Candidate (State)

Meadow checkermallow is a state Candidate species, but it has no special status at the federal level. This upright perennial herbaceous wildflower grows 20–71 inches tall and blooms between May and July (Young-Matthews 2012). Meadow checkermallow is endemic to the Willamette Valley in Oregon where it grows in meadows and prairies, along roadsides and fencerows, and at the edges of woodlands, wetlands and riparian areas. The primary threats to the species are herbicide use, habitat loss to agricultural and urban development, displacement by invasive weeds, and encroachment by trees and shrubs (Young-Matthews 2012).

Meadow checkermallow was not observed during Project surveys, and the Oregon Flora Project (2023) atlas and ORBIC (2023b) do not include reported observations within the Analysis Area. As there was some potential for suitable habitat within the Analysis Area and the species is known from neighboring counties, it was included as a species with the potential to occur. However, as the species was not observed during Project surveys, it is not expected to occur within the Analysis Area and is not expected to be adversely affected by the Project.

3.3.9 Bristly-stemmed Sidalcea

No Status (Federal); Candidate (State)

Bristly-stemmed sidalcea is a state Candidate species. This large perennial wildflower grows 28–51 inches in height and blooms as early as late April until as late as early November, with the dominant blooming period occurring June–August (Oregon Flora Project 2023). Bristly-stemmed sidalcea is endemic to western Oregon and Washington, occurring in the Oregon Coast Range, coastal areas,

and southern Washington. The species grows in open meadows, grasslands, balds, coastal bluffs, and mountain peaks (Oregon Flora Project 2023). Threats to bristly-stemmed sidalcea include plant succession that results in the conversion of meadow habitat to closed-canopy forest or overtopping shrubs, invasive plant infestation, off-highway vehicles, urban development, possible in-breeding depression due to barriers to gene flow, seed predation by weevils, possible loss of pollinators, and the effects of climate change (USFS and BLM 2011).

Bristly-stemmed sidalcea was not observed during Project surveys and there are no known occurrences within the Analysis Area. As there was some potential for suitable habitat within the Analysis Area and the species is known from neighboring counties, it was included as a species with the potential to occur. In 2023, surveys of additional areas were conducted in September, outside of the bristly-stemmed sidalcea blooming period. Potentially suitable habitat for bristly-stemmed sidalcea was identified. These additional areas will be surveyed in full prior to construction and if any bristly-stemmed sidalcea are identified they will be avoided. Therefore, the species is not expected to be adversely affected by the Project.

3.3.10 Nelson's Sidalcea

Delisted (Federal); Threatened (State)

Nelson's sidalcea was delisted from the ESA in 2023 due to recovery but remains listed as Threatened at the state level. This erect perennial wildflower grows 24–39 inches in height and blooms between late May and September (Oregon Flora Project 2023). Nelson's sidalcea is endemic to Oregon and Washington, occurring in the Willamette Valley and occasionally in the Coast Range in Oregon and north to Lewis County in central Washington. The species grows in relatively open areas on damp soil with sites including meadows, wet prairie remnants, fencerows, roadsides, deciduous forest edges, and occasionally Oregon ash wetlands (Oregon Flora Project 2023). Primary threats to Nelson's sidalcea include habitat loss due to agricultural and urban development, ecological succession resulting in the encroachment of trees and woody shrubs into open prairie habitats, and exotic weed invasions (ODA 2023, USFWS 2010). Additional threats include seed predation by weevils, possible inbreeding depression due to small population sizes and habitat fragmentation, and interspecific hybridization (ODA 2023, USFWS 2010).

Nelson's sidalcea was not observed during Project surveys and there are no known occurrences within the Analysis Area. As there was some potential for suitable habitat within the Analysis Area and the species is known from neighboring counties, it was included as a species with the potential to occur. In 2023, surveys of additional areas were conducted in September, outside of the Nelson's sidalcea blooming period. Potentially suitable habitat for Nelson's sidalcea was identified. These additional areas will be surveyed in full prior to construction and if any Nelson's sidalcea are identified they will be avoided. Therefore, the species is not expected to be adversely affected by the Project.

3.3.11 Oregon *Sullivantia*

Species of Concern (Federal); Candidate (State)

Oregon *sullivantia* is a state Candidate species and a federal Species of Concern. This delicate perennial wildflower typically grows 2–8 inches in height and blooms between late May and August (Oregon Flora Project 2023). Oregon *sullivantia* is endemic to Oregon and Washington, occurring in the Columbia River Gorge, northern Willamette Valley and Cascade Mountains in Oregon and ranging into Skamania County in southern Washington. The species grows on moist, shaded cliffs, especially near waterfalls with the surrounding forest dominated by Douglas fir (Oregon Flora Project 2023). Primary threats to Oregon *sullivantia* include collecting, recreational rock climbing, and significant changes in hydrology (Camp and Gamon 2011).

Oregon *sullivantia* was not observed during Project surveys. Because the Oregon Flora Project (2023) atlas and ORBIC (2023b) include Columbia County within the species' known range and there was some potential for suitable habitat within the Analysis Area, it was included as a species with the potential to occur. However, as the species was not observed during Project surveys, it is not expected to occur within the Analysis Area and is not expected to be adversely affected by the Project.

4.0 Avoidance and Mitigation – OAR 345-021-0010(1)(q)(C)

(C) For each species identified under (A), a description of measures proposed by the applicant, if any, to avoid or reduce adverse impact;

In accordance with OAR 345-021-0010(1)(q)(C), this section provides a description of measures proposed to avoid and minimize adverse effects to species listed in Table Q-1 and their critical habitat. Only species for which potential adverse impacts are anticipated are included.

4.1 Wildlife

4.1.1 Fish

4.1.1.1 Avoidance

Coho salmon and Chinook salmon occur in Lindgren Creek within the Analysis Area. Stream impacts will be avoided via the use of HDD to bore beneath the streambed, thereby avoiding impacts to fish species.

4.1.1.2 Minimization

The following BMPs will be used to minimize the risk of impacts to fish species:

- NWN has developed an Inadvertent Return Response Plan (Exhibit H) detailing procedures to identify and respond to an inadvertent release during HDD.

- NWN will minimize the use of herbicides to the extent practicable including avoiding their use in the vicinity of sensitive environments or species. If use of herbicides are required to control the growth of vegetation in the pipeline corridor, NWN will comply with all applicable federal and state regulations.

4.2 Plants

As discussed, in 2023 surveys of additional areas were conducted in September, outside of target species blooming periods. Areas were characterized as having potentially suitable habitat for a number of species. These additional areas will be surveyed in full prior to construction and any identified locations will be avoided.

4.2.1 *Howell's Montia*

4.2.1.1 *Avoidance*

The locations of Howell's montia within the Analysis Area were identified during Project surveys. Locations of rare plants will be avoided during construction.

4.2.1.2 *Minimization*

The following BMPs will be used to minimize the risk of impacts to any unidentified individuals:

- Prior to initiation of construction, a botanist will survey the full extent of the area where Howell's montia was identified during the 2022 botanical survey, as well as a 250-foot buffer, to identify any new plants.
- Areas with Howell's montia will be flagged with a 250-foot buffer, and those that occur in areas where proposed permanent or temporary construction activities will occur will be protected using construction safety fencing or a similar visual and physical barrier.

5.0 Protection and Conservation Program Compliance/Impacts – OAR 345-021-0010(1)(q)(D)

(D) For each plant species identified under (A), a description of how the proposed facility, including any mitigation measures, complies with the protection and conservation program, if any, that the Oregon Department of Agriculture has adopted under ORS 564.105(3);

There are no species with the potential to occur within 0.5 miles of the Project for which ODA is running a recovery program. As a result, the Project is not likely to impact any of ODA's recovery efforts, nor is the Project likely to cause a significant reduction in the likelihood of survival or recovery of plants for which ODA has developed a protection or conservation program under ORS 564.105(3).

6.0 Potential Impacts to Plants, Including Mitigation Measures – OAR 345-021-0010(1)(q)(E)

(E) For each plant species identified under paragraph (A), if the Oregon Department of Agriculture has not adopted a protection and conservation program under ORS 564.105(3), a description of significant potential impacts of the proposed facility on the continued existence of the species and on the critical habitat of such species and evidence that the proposed facility, including any mitigation measures, is not likely to cause a significant reduction in the likelihood of survival or recovery of the species;

In compliance with OAR 345-021-0010(1)(q)(E), this section describes, for plant species listed in Table Q-1 that ODA has not adopted a protection and conservation program under ORS 564.105(3), the significant potential impacts of the Project on their continued existence and on their critical habitat. This section also provides evidence, considering mitigation measures, that the Project is not likely to cause a significant reduction in their likelihood of survival or recovery.

6.1 Species Found Not to Occur

Ten species listed in Table Q-1 are not known to occur within the Analysis Area. Based on distribution information reviewed during the desktop analysis and in preparation for field work, they were considered to have the potential to occur; however, these species were not observed during surveys for the Project. Potentially suitable habitat was identified for some of these species during a botanical season conducted outside of the blooming period. A full survey will be conducted prior to construction and the species will be avoided if found. For now, we assume they are not present in the Analysis Area. As none of these species are known to occur within the Analysis Area, the Project will not cause a significant reduction in the likelihood of survival or recovery of these species:

- Tall bugbane;
- Willamette Valley larkspur;
- Peacock larkspur;
- Coast Range fawn-lily;
- Queen-of-the-forest;
- Saddle Mountain saxifrage;
- Bristly-stemmed sidalcea;
- Nelson's sidalcea;
- Meadow checkermallow; and
- Oregon sullivantia.

6.2 Species Found or Previously Known to Occur

Howell's montia, which was observed within the Site Boundary during Project surveys, is not expected to be adversely affected by the Project because the Project will avoid plants during construction. Furthermore, this species is a state candidate for listing, and receives no formal protection under the Oregon or federal ESA.

7.0 Potential Impacts to Animals, Including Mitigation Measures – OAR 345-021-0010(1)(q)(F)

(F) For each animal species identified under (A), a description of significant potential impacts of the proposed facility on the continued existence of such species and on the critical habitat of such species and evidence that the proposed facility, including any mitigation measures, is not likely to cause a significant reduction in the likelihood of survival or recovery of the species; and

In compliance with OAR 345-021-0010(1)(q)(F), this section describes the significant potential impacts of the Project on the continued existence and on the critical habitat of all animal species listed in Table Q-1, and provides evidence, taking into account mitigation measures, that the Project is not likely to cause a significant reduction in their likelihood of survival or recovery.

No impacts are anticipated to animal species identified in Table Q-1 because the Project will use HDD technology to completely avoid impacts to streams.

8.0 Monitoring – OAR 345-021-0010(1)(q)(G)

(G) The applicant's proposed monitoring program, if any, for impacts to threatened and endangered species.

No monitoring is proposed as no impacts to threatened and endangered species are not anticipated.

9.0 References

Burke Museum Herbarium. 2022a. Burke Herbarium Image Collection - Species descriptions accessed online 2022 by Erin Colclazier of Hamer Environmental. <https://burkeherbarium.org/imagecollection/browse.php?Classification=Vascular%20Plants>.

Burke Museum Herbarium. 2022b. Authors B. Legler and D. Giblin. Burke Herbarium Image Collection – *Montia howellii* Species Description. Accessed Online 2022 by Erin Colclazier of Hamer Environmental.

- <https://biology.burke.washington.edu/herbarium/imagecollection/taxon.php?Taxon=Montia%20howellii>.
- Camp, P. and J.G. Gamon (Eds.). 2011. Field Guide to the Rare Plants of Washington. University of Washington Press.
- Center for Plant Conservation. 2020. CPC National Plant Collection Profile: *Erythronium elegans*. <https://saveplants.org/plant-profile/1839/Erythronium-elegans/Coast-Range-Fawnlily/>.
- Gisler, S.D. 2004. Developing biogeographically based population introduction protocols for at-risk Willamette Valley plant species. Report to US Fish and Wildlife Service, Portland, Oregon. Native Plant Conservation Program, Oregon Department of Agriculture, Salem, Oregon.
- Google Earth. 2023. Imagery Date: 6/1/2023.
- Kaye, T.N. and J. Cramer. 2003. Direct seeding or transplanting: the cost of restoring populations of Kincaid's lupine (Oregon). *Ecological Restoration* 21:224–225.
- Meinke, R.J. 1982. Threatened and Endangered Vascular Plants of Oregon: An Illustrated Guide. U.S. Fish and Wildlife Service, Region 1, Portland, Oregon. 326 pp.
- NatureServe. 2023. NatureServe Network Biodiversity Location Data accessed through NatureServe Explorer [web application]. NatureServe, Arlington, Virginia. <https://explorer.natureserve.org/>.
- NOAA Fisheries (National Marine Fisheries Service). 2022a. National ESA Critical Habitat Mapper [web application]. Accessed August 21, 2023. <https://www.fisheries.noaa.gov/resource/map/national-esa-critical-habitat-mapper>.
- NOAA Fisheries. 2022b. 2022 5-Year Review: Summary and Evaluation of Lower Columbia River Chinook Salmon, Columbia River Chum Salmon, Lower Columbia River Coho Salmon, Lower Columbia River Steelhead. <https://www.fisheries.noaa.gov/resource/document/2022-5-year-review-summary-evaluation-lower-columbia-river-chinook-salmon>.
- NOAA Fisheries. 2023. Species information, maps, and GIS data website queries. Accessed August 22, 2023. http://www.westcoast.fisheries.noaa.gov/maps_data/maps_and_gis_data.html.
- ODA (Oregon Department of Agriculture). 2022. About the Plants. Plant Conservation Program. <https://www.oregon.gov/oda/programs/PlantConservation/Pages/AboutPlants.aspx>.
- ODA. 2023. Nelson's checkermallow species profile. <http://www.oregon.gov/ODA/shared/Documents/Publications/PlantConservation/SidalceaNelsonianaProfile.pdf>.
- ODFW (Oregon Department of Fish and Wildlife). 2007. Oregon Coast Coho Conservation Plan for the State of Oregon. March 16, 2007. https://www.dfw.state.or.us/fish/CRP/docs/coastal_coho/final/Coho_Plan.pdf


- ODFW. 2010. Lower Columbia River Conservation and Recovery Plan for Oregon Populations of Salmon and Steelhead. August 7, 2010. https://www.dfw.state.or.us/fish/CRP/docs/lower-columbia/OR_LCR_Plan%20-%20Aug_6_2010_Final.pdf
- ODFW. 2017. 2017 ODFW Western Oregon Deer and Elk Habitat. <https://www.dfw.state.or.us/habitat/mitigation/Final%202017%20ODFW%20WO%20Deer%20and%20Elk%20Habitat%20Mapping%20Rationale%20-%20April%202017.pdf>
- ODFW 2023a. White-tailed deer species profile. Accessed August 22, 2023. http://www.dfw.state.or.us/species/mammals/hoofed_mammals.asp.
- ODFW. 2023b. Species information, maps, and GIS data website queries. <https://www.dfw.state.or.us/wildlife/diversity/species/index.asp>.
- ODFW. 2023c. Oregon guidelines for timing of in-water work to protect fish and wildlife resources (April 2023). <https://www.dfw.state.or.us/lands/inwater/2023%20Oregon%20In-Water%20Work%20Guidelines.pdf>.
- ODFW. 2023d. Natural Resources Information Management Program: Oregon fish life history timing tables. <https://nrimp.dfw.state.or.us/DataClearinghouse/default.aspx?p=202&XMLname=42654.xml> Accessed January 2024.
- ODFW. 2024. Oregon Fish Habitat and Distribution Barriers. https://nrimp.dfw.state.or.us/FHD_FPB_Viewer/index.html Accessed January 2024.
- ORBIC (Oregon Biodiversity Information Center). 2019. ORBIC Rare, Threatened and Endangered species of Oregon. Oregon Biodiversity Information Center. <https://inr.oregonstate.edu/sites/inr.oregonstate.edu/files/2019-rte-book.pdf>.
- ORBIC. 2023a. ORBIC Rare, Threatened and Endangered species of Oregon. Oregon Biodiversity Information Center. In Prep. <https://inr.oregonstate.edu/orbic/rare-species/rare-species-oregon-publications>.
- ORBIC. 2023b. GIS data provided in response to data request for GIS data pertaining to listed, endangered, threatened, or special concern species occurring within Clatsop, Columbia, Tillamook, and Washington counties in Oregon. Oregon Biodiversity Information Center, Portland, Oregon.
- Oregon Flora Project. 2023. Taxa review of state and federal listed vascular plant species distribution and habitats. OregonFlora, Oregon State University, Department of Botany and Plant Pathology, Corvallis, Oregon. <https://oregonflora.org/>.
- Polfus, J.L. 2011. Literature review and synthesis on the effects of residential development on ungulate winter range in the Rocky Mountain West. Report prepared for Montana Fish, Wildlife and Parks. Helena, MT.

- Renner, M.A., J.Regan, and M. Colosio. 2012. Response of *Montia howellii* (Howell's montia) to Road Management in California pp 303–312 in Proceedings of coast redwood forests in a changing California: A symposium for scientists and managers, PSW-GTR-238.
- StreamNet. 2024. StreamNet Mapper [web application]. StreamNet Program.
<https://www.streamnet.org/home/data-maps/sn-mapper/>. Accessed January 2024.
- USFS (U.S. Department of Agriculture, Forest Service) and BLM (U.S. Department of Interior, Bureau of Land Management). 2011. Conservation assessment for bristly-stemmed checker-mallow (*Sidalcea hirtipes*). USDA Forest Service Region 6 and USDI Bureau of Land Management, Oregon and Washington.
- USFWS (U.S. Fish and Wildlife Service). 1983. Columbian white-tailed deer recovery plan. Accessed August 22, 2023. https://ecos.fws.gov/docs/recovery_plan/830614.pdf.
- USFWS. 2010. Recovery Plan for the Prairie Species of Western Oregon and Southwestern Washington. U.S. Fish and Wildlife Service, Portland, Oregon. xi + 241 pp.
- USFWS. 2016. Endangered and Threatened Wildlife and Plants; Reclassifying the Columbia River Distinct Population Segment of the Columbian White-Tailed Deer as Threatened With a Rule Under Section 4(d) of the Act. Accessed August 22, 2023.
<https://ecos.fws.gov/ecp/species/154>.
- USFWS. 2023. IPaC [web application]. Accessed August 22, 2023. <https://ipac.ecosphere.fws.gov/>.
- WNHP (Washington Natural Heritage Program). 2014. Field guide to selected rare plants of Washington. Online. <http://www1.dnr.wa.gov/nhp/refdesk/fguide/pdf/ciel.pdf>.
- WNHP. 2022. Washington Vascular Plant Species of Special Concern by County. Washington Natural Heritage Program, Washington Department of Natural Resources. Olympia, WA.
https://www.dnr.wa.gov/publications/amp_nh_county_plants.pdf?gn8v6t.

Figures

(Figure Q-3 and Figure Q-4 are confidential and submitted under a separate cover.)

This page intentionally left blank



I:\cscs\706\gis\1\CES\Projects\PD\X\NW_NaturalMistResiliency\Maps_p\RA\13\Exhibit_Q\NW_N_MistResiliency_ExhibitQ_20240214.aprx


1:40,000
 WGS 1984 UTM Zone 10N
0
1 Miles
NOT FOR CONSTRUCTION


Mist Resiliency Project

**Figure Q-1
Threatened and Endangered
Species Analysis Area**

COLUMBIA COUNTY, OREGON

- Site Boundary
- County Boundary
- US Highway
- State Highway





\\cass706\gis\1\CES\Projects\PDX\NW_Natural\MistResiliency\Maps_p\RA13\Exhibit_Q\NW_N_MistResiliency_ExhibitQ_20240214.aprx

1:40,000 WGS 1984 UTM Zone 10N 0 1 Miles NOT FOR CONSTRUCTION

Mist Resiliency Project

**Figure Q-2
Surveyed Area:
Threatened and
Endangered Plant Species**

COLUMBIA COUNTY, OREGON

-  Site Boundary
-  Botanical Survey Area
-  City/Town
-  County Boundary
-  US Highway
-  State Highway

