Quick Reference 1-9

TABLE 1-6. Oregon Rates of Low Birthweight, and Measures of Prenatal Care, 1980-2001

| Year | Low Birthweight | First Trimester | No Care | Inadequate Care | Third Trimester Care | Less than Five Visits |
|------|--------------------|--------------------|---------|--------------------|----------------------------|--------------------------|
| 1980 | 50.4 | 780.8 | 5.5 | 58.0 | 35.2 | 41.4 |
| 1981 | 48.5 | 775.6 | 8.9 | 63.1 | 38.6 | 43.0 |
| 1982 | 49.2 | 769.3 | 11.2 | 70.3 | 41.0 | 48.0 |
| 1983 | 50.0 | 775.3 | 11.3 | 66.5 | 38.5 | 44.9 |
| 1984 | 51.5 | 771.5 | 11.0 | 68.2 | 41.1 | 46.2 |
| 1985 | 51.3 | 752.0 | 12.1 | 72.9 | 43.7 | 47.5 |
| 1986 | 51.3 | 738.7 | 11.7 | 83.3 | 52.1 | 54.6 |
| 1987 | 54.0 | 736.8 | 16.5 | 86.2 | 50.3 | 58.5 |
| 1988 | 52.6 | 738.8 | 13.8 | 83.6 | 49.9 | 54.7 |
| 1989 | 52.2 | 750.7 | 12.0 | 73.2 | 42.9 | 48.7 |
| 1990 | 50.1 | 757.1 | 10.7 | 70.0 | 43.4 | 45.1 |
| 1991 | 49.2 | 768.2 | 8.7 | 61.0 | 37.4 | 38.6 |
| 1992 | 51.8 | 787.0 | 8.2 | 52.6 | 31.4 | 34.0 |
| 1993 | 52.5 | 794.6 | 7.6 | 51.7 | 30.4 | 33.8 |
| 1994 | 53.0 | 790.9 | 8.5 | 57.8 | 34.3 | 36.4 |
| 1995 | 54.9 | 787.7 | 8.6 | 58.4 | 34.7 | 38.2 |
| 1996 | 53.5 | 799.3 | 7.1 | 53.7 | 31.7 | 34.8 |
| 1997 | 55.0 | 811.2 | 6.7 | 50.0 | 29.6 | 32.3 |
| 1998 | 53.7 | 807.2 | 7.2 | 53.5 | 30.7 | 35.3 |
| 1999 | 53.9 | 809.9 | 7.3 | 53.7 | 29.6 | 35.7 |
| 2000 | 56.6 | 812.8 | 8.5 | 55.9 | 29.8 | 36.6 |
| 2001 | 55.6 | 815.2 | 8.0 | 50.5 | 28.7 | 33.1 |

Inadequate prenatal care is defined as care that began in the third trimester or consisted of less than five visits. All rates are per 1,000 live births. Rates are calculated excluding missing and unknown values.