

TABLE 2-24. Selected Medical or Health Characteristics by Mother's Age (Percents), Oregon Resident Births, 2010

| Characteristic | Total Births ¹ | Age of Mother | | | | | | | |
|--|---------------------------|---------------|-------|--------|--------|--------|-------|-------|------|
| | | <15 | 15-19 | 20-24 | 25-29 | 30-34 | 35-39 | 40-44 | 45+ |
| All Births - Mother | | | | | | | | | |
| Total Births | 45,596 | 27 | 3,511 | 10,325 | 13,381 | 11,480 | 5,580 | 1,202 | 90 |
| First Trimester Care | 73.1 | 46.2 | 57.9 | 65.4 | 75.0 | 79.6 | 78.5 | 74.5 | 76.7 |
| Inadequate Care ² | 5.5 | 19.2 | 9.5 | 7.6 | 5.2 | 3.6 | 3.8 | 4.0 | 4.7 |
| No Prenatal Care | 0.6 | 3.8 | 0.8 | 0.8 | 0.7 | 0.4 | 0.5 | 0.2 | 1.2 |
| Out-Of-Hospital Birth .. | 3.4 | 3.7 | 0.8 | 1.9 | 3.6 | 4.6 | 4.6 | 4.7 | 1.1 |
| Primary Cesarean | 18.2 | 22.2 | 16.7 | 17.2 | 16.9 | 18.9 | 20.9 | 26.0 | 41.1 |
| Repeat Cesarean | 11.1 | – | 2.0 | 8.3 | 10.9 | 13.3 | 16.5 | 17.9 | 24.4 |
| Multiple Births | 3.4 | – | 1.0 | 2.6 | 2.8 | 4.0 | 5.1 | 7.6 | 25.6 |
| Tobacco Use | 11.3 | 3.7 | 18.5 | 18.5 | 11.4 | 6.4 | 5.1 | 5.0 | 5.6 |
| Overweight/Obese ³ | 48.3 | – | 37.3 | 49.1 | 50.0 | 48.4 | 48.4 | 52.2 | 56.8 |
| All Births - Infant | | | | | | | | | |
| Preterm Births ⁴ | 7.9 | 7.4 | 8.4 | 7.8 | 7.4 | 7.3 | 9.3 | 11.8 | 27.8 |
| Very Low Birthweight ⁵ | 1.1 | – | 1.3 | 1.1 | 1.0 | 0.8 | 1.3 | 1.6 | 6.7 |
| Low Birthweight ⁶ | 6.3 | – | 7.7 | 6.7 | 5.5 | 5.7 | 7.2 | 7.9 | 21.1 |
| Fetal Macrosomia ⁷ | 10.4 | – | 6.5 | 8.6 | 10.7 | 12.0 | 11.8 | 12.1 | 5.6 |
| 5 Minute Apgar < 7 | 2.7 | 14.8 | 3.6 | 2.8 | 2.5 | 2.4 | 2.7 | 3.2 | 7.9 |
| Mothers Born in the U.S. | | | | | | | | | |
| Total Births | 35,663 | 23 | 2,992 | 8,443 | 10,558 | 8,776 | 3,969 | 853 | 49 |
| First Trimester Care | 75.3 | 50.0 | 60.0 | 67.0 | 77.9 | 82.4 | 81.1 | 78.6 | 83.7 |
| Inadequate Care ² | 5.0 | 18.2 | 9.0 | 7.0 | 4.6 | 3.2 | 3.1 | 3.1 | 4.2 |
| No Prenatal Care | 0.6 | 4.5 | 0.9 | 0.8 | 0.5 | 0.4 | 0.5 | 0.1 | – |
| Out-Of-Hospital Birth .. | 4.0 | 4.3 | 0.7 | 2.1 | 4.2 | 5.5 | 5.7 | 6.2 | – |
| Primary Cesarean | 18.7 | 21.7 | 16.9 | 17.8 | 17.6 | 19.1 | 21.6 | 25.8 | 36.7 |
| Repeat Cesarean | 10.6 | – | 1.9 | 8.2 | 10.6 | 12.6 | 16.1 | 17.5 | 26.5 |
| Multiple Births | 3.5 | – | 1.0 | 2.7 | 3.2 | 4.0 | 5.8 | 7.6 | 28.6 |
| Tobacco Use | 14.2 | 4.3 | 21.6 | 22.2 | 14.1 | 8.1 | 6.8 | 6.4 | 10.2 |
| Overweight/Obese ³ | 48.4 | – | 37.7 | 50.3 | 50.5 | 48.1 | 47.2 | 50.4 | 63.3 |
| Infants of Mothers Born in the U.S. | | | | | | | | | |
| Preterm Births ⁴ | 7.9 | 4.3 | 8.4 | 8.0 | 7.5 | 7.3 | 9.1 | 11.0 | 32.7 |
| Very Low Birthweight ⁵ | 1.0 | – | 1.1 | 1.1 | 1.0 | 0.7 | 1.2 | 0.7 | 6.1 |
| Low Birthweight ⁶ | 6.3 | – | 7.7 | 6.9 | 5.8 | 5.5 | 7.1 | 6.8 | 20.4 |
| Fetal Macrosomia ⁷ | 10.7 | – | 6.7 | 8.7 | 11.1 | 12.6 | 12.4 | 13.1 | 4.1 |
| 5 Minute Apgar < 7 | 2.8 | 13.0 | 3.6 | 2.9 | 2.7 | 2.5 | 2.8 | 3.1 | 6.1 |

– Quantity is zero.
See footnotes at end of table.

TABLE 2-24. Selected Medical or Health Characteristics by Mother's Age (Percents), Oregon Resident Births, 2010 (Continued)

| Characteristic | Total Births | Age of Mother | | | | | | | |
|---|--------------|---------------|-------|-------|-------|-------|-------|-------|------|
| | | <15 | 15-19 | 20-24 | 25-29 | 30-34 | 35-39 | 40-44 | 45+ |
| Mothers Born Outside the U.S. | | | | | | | | | |
| Total Births | 9,933 | 4 | 519 | 1,882 | 2,823 | 2,704 | 1,611 | 349 | 41 |
| First Trimester Care | 65.2 | 25.0 | 46.0 | 58.2 | 64.4 | 70.7 | 72.2 | 64.5 | 68.3 |
| Inadequate Care ² | 7.1 | 25.0 | 12.0 | 10.0 | 7.4 | 4.9 | 5.3 | 6.2 | 5.3 |
| No Prenatal Care | 0.8 | – | 0.8 | 0.9 | 1.0 | 0.5 | 0.6 | 0.3 | 2.6 |
| Out-Of-Hospital Birth .. | 1.3 | – | 1.0 | 0.9 | 1.2 | 1.4 | 1.9 | 0.9 | 2.4 |
| Primary Cesarean | 16.8 | 25.0 | 15.6 | 14.6 | 14.1 | 18.2 | 19.2 | 26.6 | 46.3 |
| Repeat Cesarean | 13.1 | – | 2.3 | 8.7 | 12.2 | 15.5 | 17.6 | 18.9 | 22.0 |
| Multiple Births | 2.8 | – | 0.8 | 2.2 | 1.5 | 3.7 | 3.4 | 7.4 | 22.0 |
| Tobacco Use | 1.2 | – | 1.2 | 2.1 | 1.1 | 1.0 | 0.8 | 1.5 | – |
| Overweight/Obese ³ | 47.9 | – | 35.1 | 43.7 | 48.4 | 49.5 | 51.4 | 56.7 | 48.7 |
| Infants of Mothers Born Outside the U.S. | | | | | | | | | |
| Preterm Births ⁴ | 7.8 | 25.0 | 8.1 | 6.9 | 6.7 | 7.2 | 9.7 | 13.8 | 22.0 |
| Very Low Birthweight ⁵ | 1.4 | – | 1.9 | 1.4 | 0.9 | 1.2 | 1.6 | 3.7 | 7.3 |
| Low Birthweight ⁶ | 6.3 | – | 7.9 | 6.1 | 4.6 | 6.2 | 7.6 | 10.6 | 22.0 |
| Fetal Macrosomia ⁷ | 9.1 | – | 5.8 | 7.9 | 8.8 | 10.0 | 10.2 | 9.7 | 7.3 |
| 5 Minute Apgar < 7 | 2.3 | 25.0 | 3.7 | 2.1 | 1.9 | 2.3 | 2.3 | 3.7 | 10.0 |

- Quantity is zero.
- 1 There were no births with unknown age of mother.
- 2 Less than 5 prenatal visits or care began in the third trimester.
- 3 Body Mass Index of greater than 25.0 kg/m².
- 4 Born prior to 37 completed weeks of gestation.
- 5 Birthweight of less than 1,500 grams (3 lb 4 oz).
- 6 Birthweight of less than 2,500 grams (5 lb 8 oz).
- 7 Birthweight of more than 4,000 grams (8 lb 13 oz).

NOTE: Rates and percentages are calculated excluding missing and unknown values.