

TABLE 7-18. Neonatal, Postneonatal, and Infant Death Rates by Mother's Risk Factors, Oregon Residents, Birth Cohort 2000-2002

Risk Factor	Neonatal ¹		Post-Neonatal ²		Infant ³	
	No.	Rate	No.	Rate	No.	Rate
Total	489	3.6	262	1.9	751	5.5
Marital Status						
Married	309	3.3	117	1.2	426	4.5
Unmarried	180	4.3	145	3.5	325	7.8
Mother's Age						
10-14	1	–	1	–	2	–
15-19	59	4.1	56	3.9	115	8.0
20-24	133	3.6	81	2.2	214	5.9
25-29	124	3.3	66	1.7	190	5.0
30-34	96	3.2	29	1.0	125	4.1
35-39	61	4.4	24	1.7	85	6.1
40-44	11	3.6	3	–	14	4.6
45+	2	–	1	–	3	–
Non-Hispanic						
White	362	3.6	196	1.9	558	5.5
African American	14	5.0	14	5.0	28	10.0
American Indian	9	4.4	5	2.5	14	6.9
Asian ⁴	18	2.6	12	1.8	30	4.4
Total Hispanic	84	3.6	33	1.4	117	5.0
Mother's Education						
8 th Grade or Less	25	2.8	16	1.8	41	4.6
Some High School	83	4.5	78	4.2	161	8.7
HS diploma/GED	164	3.8	82	1.9	246	5.7
More than High School	185	2.9	77	1.2	262	4.1
Start of Prenatal Care						
1 st Trimester	381	3.4	181	1.6	562	5.1
2 nd Trimester	69	3.4	57	2.8	126	6.3
3 rd Trimester	10	2.5	18	4.6	28	7.1
No Care	29	20.9	6	4.3	35	25.3
Tobacco Use						
Yes	74	4.2	79	4.5	153	8.8
No	387	3.3	178	1.5	565	4.8
Alcohol Use						
Yes	7	4.2	4	–	11	6.6
No	453	3.4	248	1.9	701	5.3
Multiple Birth						
Yes	84	21.4	12	3.1	96	24.4
No	405	3.1	250	1.9	655	4.9

– Quantity is zero or rate is based on fewer than five occurrences.

NOTE: Because of unreported items, the sum of all categories may not equal the total.

All rates per 1,000 live births.

¹ Neonatal deaths include infant deaths of less than 28 days.

² Postneonatal deaths occur from day 28 through 364 after birth.

³ Infant death is the death of a child prior to its first birthday.

⁴ Includes Chinese, Japanese, Filipino, and Other Asian & Pacific Islander.