

TABLE 6-2. Leading causes of death for males and females by rank order, number, rate, percent, and median age at death, Oregon residents, 2013

Cause of death in rank order	Rank	No.	Rate ¹	Pct.	Median age
Males					
Total		17,171	886.8	100.0	75
Malignant neoplasms	1	4,115	212.5	24.0	72
Diseases of the heart	2	3,509	181.2	20.4	79
Unintended injuries	3	996	51.4	5.8	57
Chronic lower respiratory disease	4	965	49.8	5.6	77
Cerebrovascular disease	5	756	39.0	4.4	81
Diabetes mellitus	6	634	32.7	3.7	73
Suicide	7	536	27.7	3.1	49
Alcohol-induced	8	516	26.6	3.0	58
Alzheimer's disease	9	401	20.7	2.3	87
Hypertension & hyp. renal disease	10	255	13.2	1.5	77
Parkinson's disease	11	235	12.1	1.4	82
Influenza & pneumonia	12	229	11.8	1.3	84
Viral hepatitis	13	167	8.6	1.0	59
Nephritis, nephrotic syndrome, etc.	14	156	8.1	0.9	82
Neoplasms not known to be malignant	15	132	6.8	0.8	77
Septicemia	16	118	6.1	0.7	75
Aortic aneurysm	17	92	4.8	0.5	77
Pneumonitis due to solids & liquids	18	84	4.3	0.5	83
Amyotrophic lateral sclerosis	19	75	3.9	0.4	67
Congenital malformations	20	74	3.8	0.4	36
Females					
Total		16,760	845.3	100.0	82
Malignant neoplasms	1	3,683	185.8	22.0	73
Diseases of the heart	2	2,988	150.7	17.8	86
Chronic lower respiratory disease	3	1,060	53.5	6.3	78
Cerebrovascular disease	4	1,013	51.1	6.0	85
Alzheimer's disease	5	910	45.9	5.4	88
Unintended injuries	6	736	37.1	4.4	76
Diabetes mellitus	7	477	24.1	2.8	78
Influenza & pneumonia	8	272	13.7	1.6	84
Hypertension & hyp. renal disease	9	268	13.5	1.6	88
Alcohol-induced	10	197	9.9	1.2	55
Nephritis, nephrotic syndrome, etc.	11	171	8.6	1.0	82
Suicide	12	161	8.1	1.0	50
Parkinson's disease	13	159	8.0	0.9	85
Neoplasms not known to be malignant	14	116	5.9	0.7	82
Septicemia	15	104	5.2	0.6	79
Pneumonitis due to solids & liquids	16	72	3.6	0.4	85
Viral hepatitis	17	67	3.4	0.4	58
Congenital malformations	18	66	3.3	0.4	36
Perinatal conditions	19	65	3.3	0.4	0
Amyotrophic lateral sclerosis	20	64	3.2	0.4	70

¹ All rates per 100,000 population.