

**TABLE 6-20. Tobacco-linked deaths by cause of death, Oregon residents, 2013**

Selected causes of death (and their ICD-10 codes)	Total	Linked <sup>1</sup>		Not linked		Unknown	
		Number	Percent	Number	Percent	Number	Percent
Total <sup>2</sup> .....	33,931	7,436	21.9	18,672	55.0	7,823	23.1
Malignant neoplasms .....	3,522	1,892	53.7	1,042	29.6	588	16.7
Oral cavity, lip, pharynx (C00.0-C14.8) .....	118	58	49.2	39	33.1	21	17.8
Esophagus (C15) .....	213	72	33.8	75	35.2	66	31.0
Stomach (C16) .....	109	13	11.9	75	68.8	21	19.3
Pancreas (C25) .....	462	62	13.4	293	63.4	107	23.2
Larynx (C32) .....	38	35	92.1	–	–	3	7.9
Lung, bronchi, and trachea (C33-C34) .....	2,001	1,557	77.8	187	9.3	257	12.8
Cervix uteri (C53) .....	41	4	9.8	34	82.9	3	7.3
Kidney, other urinary tract (C64-C65) .....	182	16	8.8	121	66.5	45	24.7
Urinary bladder (C67) .....	206	66	32.0	87	42.2	53	25.7
Acute myeloid leukemia (C92.0) .....	152	9	5.9	131	86.2	12	7.9
Cardiovascular disease .....	8,357	1,891	22.6	4,142	49.6	2,324	27.8
Ischemic heart disease (I20-I25) .....	3,421	1,116	32.6	1,394	40.7	911	26.6
Other heart disease (I00-I09, I26-I51) .....	2,814	401	14.3	1,647	58.5	766	27.2
Cerebrovascular disease (I60-I69) .....	1,769	244	13.8	978	55.3	547	30.9
Atherosclerosis (I70) .....	59	18	30.5	30	50.8	11	18.6
Aortic aneurysm (I71) .....	155	48	31.0	58	37.4	49	31.6
Other arterial disease (I72-I78) .....	139	64	46.0	35	25.2	40	28.8
Respiratory diseases .....	2,453	1,599	65.2	454	18.5	400	16.3
Pneumonia and influenza (J09-J18) .....	501	59	11.8	303	60.5	139	27.7
Bronchitis and emphysema (J40-J43) .....	170	139	81.8	18	10.6	13	7.6
Other chronic airways obstruction (J44) .....	1,782	1,401	78.6	133	7.5	248	13.9
Perinatal conditions <sup>3</sup> .....	68	3	4.4	63	92.6	2	2.9
Selected perinatal conditions <sup>4</sup> .....	45	1	2.2	42	93.3	2	4.4
Sudden infant death syndrome (R95) .....	23	2	8.7	21	91.3	–	–
Other causes .....	19,531	2,051	10.5	12,971	66.4	4,509	23.1

<sup>1</sup> The Oregon death certificate asks 'Did tobacco use contribute to death?' This is followed by four checkboxes: 'Yes,' 'No,' 'Probably,' and 'Unknown.' The linked category includes deaths listed as 'Yes' or 'Probably.'

<sup>2</sup> The causes of death shown in this table are those linked to tobacco use by the federal Centers for Disease Control and Prevention (CDC. Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses -- United States, 1997-2001. MMWR 2005; 54:625-628.).

<sup>3</sup> The number of infant deaths resulting from exposure to tobacco combustion products is underreported.

<sup>4</sup> The category includes the following conditions: other disorders related to short gestation and low birthweight (P07), respiratory distress of newborn (P22), congenital pneumonia (P23), neonatal aspiration syndromes (P24), and other respiratory conditions originating in the perinatal period (P25-P28).

– Quantity is zero.