

TABLE 7-18. Neonatal, postneonatal and infant death rates by mother's risk factors, Oregon residents, birth cohort 2011-2013

Risk factors	Neonatal ¹		Postneonatal ²		Infant ³	
	No.	Rate	No.	Rate	No.	Rate
Total⁴	468	3.5	211	1.6	679	5.0
Marital status						
Married	282	3.2	87	1.0	369	4.2
Unmarried	179	3.7	124	2.6	303	6.3
Age of mother						
10-14	—	—	—	—	—	—
15-19	30	3.5	23	2.7	53	6.2
20-24	92	3.2	65	2.2	157	5.4
25-29	124	3.2	63	1.6	187	4.8
30-34	122	3.3	31	0.8	153	4.2
35-39	67	3.8	23	1.3	90	5.1
40-44	28	7.3	5	1.3	33	8.7
45+	4	*	1	*	5	19.8
Non-Hispanic race						
White	310	3.3	135	1.4	445	4.8
Black	15	5.4	10	3.6	25	9.0
American Indian	12	7.4	10	6.2	22	13.6
Asian ⁵	18	2.8	10	1.6	28	4.4
Pacific Islander ⁶	4	*	1	*	5	5.7
Other & unknown	9	16.4	1	*	10	18.2
Two or more races	10	2.4	8	1.9	18	4.2
Total Hispanic	90	3.5	36	1.4	126	4.9
Education						
8th grade or less	25	4.5	6	1.1	31	5.6
Some high school	51	3.2	47	2.9	98	6.1
HS diploma/GED	100	3.3	53	1.7	153	5.0
More than HS	273	3.3	103	1.3	376	4.6
Start of prenatal care						
Any trimester	386	3.0	186	1.5	572	4.5
1st trimester	299	3.0	118	1.2	417	4.2
2nd trimester	80	3.4	56	2.4	136	5.9
3rd trimester	7	1.5	12	2.6	19	4.1
No prenatal care	19	21.1	2	*	21	23.3
Tobacco use						
Pre-pregnancy only	17	6.0	3	*	20	7.1
During pregnancy	54	3.9	62	4.4	116	8.3
No tobacco use	386	3.3	142	1.2	528	4.5
Multiple birth						
Yes	101	22.6	18	4.0	119	26.7
No	367	2.8	193	1.5	560	4.3

1 Neonatal deaths include infant deaths of less than 28 days.

2 Postneonatal deaths occur from day 28 through 364 after birth.

3 Infant death is the death of a child prior to its first birthday.

4 Due to unreported items, the sum of all categories may not equal the total.

5 Includes Asian Indian, Chinese, Filipino, Japanese, Korean, Vietnamese and other Asian.

6 Includes Guamanian, Hawaiian, Samoan and other Pacific Islander.

* Rates are not calculated when there are fewer than 5 deaths in a category.

— Quantity is zero.

NOTE: All rates per 1,000 live births.