

TABLE 7-9. Late neonatal deaths¹ by weeks of gestation and weight, Oregon residents, birth cohort 2015

Birthweight (in grams)	Total	Weeks of gestation ²								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total ³	25	–	–	5	4	4	1	8	3	–
001-349	–	–	–	–	–	–	–	–	–	–
350-499	–	–	–	–	–	–	–	–	–	–
<500	–	–	–	–	–	–	–	–	–	–
500-749	3	–	–	2	1	–	–	–	–	–
750-999	2	–	–	2	–	–	–	–	–	–
1000-1249	2	–	–	1	1	–	–	–	–	–
1250-1499	3	–	–	–	2	1	–	–	–	–
1500-1999	–	–	–	–	–	–	–	–	–	–
2000-2499	2	–	–	–	–	1	1	–	–	–
<2500	12	–	–	5	4	2	1	–	–	–
2500+	13	–	–	–	–	2	–	8	3	–
2500-2999	7	–	–	–	–	2	–	5	–	–
3000-3499	4	–	–	–	–	–	–	3	1	–
3500-3999	2	–	–	–	–	–	–	–	2	–
4000-4499	–	–	–	–	–	–	–	–	–	–
4500+	–	–	–	–	–	–	–	–	–	–

– Quantity is zero.

¹ Late neonatal deaths occur from day seven through 27 after birth.

² Clinical estimate of gestation. If clinical estimate is unknown, gestation is calculated from reported date of last menses.

³ Total includes reports with unknown birthweight.