

TABLE 6-20. Tobacco-linked deaths by cause of death, Oregon residents, 2017

Selected causes of death (and their ICD-10 codes)	Total	Linked ¹		Not linked		Unknown	
		Number	Percent	Number	Percent	Number	Percent
Total ²	36,640	7,843	21.4	20,053	54.7	8,744	23.9
Malignant neoplasms	3,670	1,830	49.9	1,159	31.6	681	18.6
Oral cavity, lip, pharynx (C00.0-C14.8)	145	68	46.9	44	30.3	33	22.8
Esophagus (C15)	249	109	43.8	88	35.3	52	20.9
Stomach (C16)	97	22	22.7	40	41.2	35	36.1
Pancreas (C25)	640	87	13.6	416	65.0	137	21.4
Larynx (C32)	27	19	70.4	2	7.4	6	22.2
Lung, bronchi, and trachea (C33-C34)	1,865	1,401	75.1	210	11.3	254	13.6
Cervix uteri (C53)	47	4	8.5	30	63.8	13	27.7
Kidney, other urinary tract (C64-C65)	190	33	17.4	103	54.2	54	28.4
Urinary bladder (C67)	254	83	32.7	93	36.6	78	30.7
Acute myeloid leukemia (C92.0)	156	4	2.6	133	85.3	19	12.2
Cardiovascular disease	9,009	2,048	22.7	4,226	46.9	2,735	30.4
Ischemic heart disease (I20-I25)	3,387	1,098	32.4	1,323	39.1	966	28.5
Other heart disease (I00-I09, I26-I51)	3,198	525	16.4	1,730	54.1	943	29.5
Cerebrovascular disease (I60-I69)	2,066	304	14.7	1,039	50.3	723	35.0
Atherosclerosis (I70)	31	12	38.7	10	32.3	9	29.0
Aortic aneurysm (I71)	179	58	32.4	64	35.8	57	31.8
Other arterial disease (I72-I78)	148	51	34.5	60	40.5	37	25.0
Respiratory diseases	2,577	1,698	65.9	482	18.7	397	15.4
Pneumonia and influenza (J09-J18)	573	83	14.5	334	58.3	156	27.2
Bronchitis and emphysema (J40-J43)	153	129	84.3	13	8.5	11	7.2
Other chronic airways obstruction (J44)	1,851	1,486	80.3	135	7.3	230	12.4
Perinatal conditions ³	67	—	—	64	95.5	3	4.5
Selected perinatal conditions ⁴	46	—	—	45	97.8	1	2.2
Sudden infant death syndrome (R95)	21	—	—	19	90.5	2	9.5
Other causes	21,317	2,267	10.6	14,122	66.2	4,928	23.1

[—] Quantity is zero.

¹ The Oregon death certificate provides four possible answers to the question, "Did tobacco use contribute to death?": yes, probably, no and unknown. The linked category includes deaths listed as yes or probably, or if a contributing cause is F17.

² The causes of death shown in this table are those linked to tobacco use by the federal Centers for Disease Control and Prevention (CDC. Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses -- United States, 1997-2001. MMWR 2005; 54:625-628.).

³ The number of infant deaths resulting from exposure to tobacco combustion products is underreported.

⁴ The category includes the following conditions: other disorders related to short gestation and low birthweight (P07), respiratory distress of newborn (P22), congenital pneumonia (P23), neonatal aspiration syndromes (P24), and other respiratory conditions originating in the perinatal period (P25-P28).